

## Solution Set #1

### Section 6.1

1. Example 1. Let  $M = \mathbf{R}^1$  and let  $A = (0,1)$ . Then every open subset  $O$  of  $A$  can be written as  $O = A \cap B$  for  $B$  some open subset of  $M$ . But since  $A$  is open in  $M$ , then  $A \cap B$  is open in  $M$ .

Example 2. Let  $M = \mathbf{R}_d$  and let  $A$  be any subset of  $M$ .

2.

		$[0,1]$	$\mathbf{R}^1$	$\mathbf{R}^2$
a)	$(1/2,1]$	open	not	not
b)	$(1/2,1)$	open	open	not
c)	$[1/2,1)$	not	not	not

3. The only subset of  $\mathbf{R}^1$  which is open in  $\mathbf{R}^2$  is the empty set.

### Section 6.2


- Suppose that  $f$  is non-constant. Then there exist  $c, d \in f(\mathbf{R}^1)$ , such that  $c \neq d$ . Wolog  $c < d$ . By corollary 6.2E  $f$  takes on every value between  $c$  and  $d$ . But the interval  $[c,d]$  is uncountable.
- Suppose such an  $f$  existed. Let  $[a,b] \subset \mathbf{R}^1$ ,  $a < b$ . Let  $[a,b]_Q = [a,b] \cap \mathbf{Q}$  and  $[a,b]_I = [a,b] \cap \mathbf{J}$ , i.e.,  $[a,b]_Q$  is the subset of  $[a,b]$  of rationals and  $[a,b]_I$  is the subset of  $[a,b]$  of irrationals. Since  $[a,b]_Q$  is a subset of  $\mathbf{Q}$ , then  $[a,b]_Q$  is countable and  $f([a,b]_Q)$  is therefore countable. By hypothesis  $f([a,b]_I)$  is a subset of  $\mathbf{Q}$  and therefore countable. Hence  $f([a,b]) = f([a,b]_Q) \cup f([a,b]_I)$ , is countable. But by corollary 6.2E  $f([a,b])$  must contain the interval between  $f(a)$  and  $f(b)$ , which is uncountable. This is a contradiction.
- Let  $A = [0,1/2)$  and  $B = [1/2,1]$ . Then since every subset of  $\mathbf{R}_d$  is open, we have both  $A$  and  $B$  are open. Together they form a non-trivial disjoint open decomposition of  $[0,1]$ . By 6.2A part (b),  $[0,1]$  is not connected.
- False. Let  $A = [0,1]$ . Let  $B = [0,1] \cup [2,3]$ . Let  $C = \mathbf{R}^1$ .
- Wolog we may suppose that  $B \setminus A$  is non-empty, i.e.,  $B$  is strictly bigger than  $A$ . Suppose to the contrary that  $B$  were disconnected. Then, there would exist a non-trivial disjoint open decomposition of  $B$ , say sets  $C$  and  $D$ . Since  $A$  is connected, we must have either  $A \subset C$  or else  $A \subset D$ . (Otherwise the sets  $A \cap C$  and  $A \cap D$  would form non-trivial disjoint open decomposition of  $A$ .) Wolog  $A \subset C$ . Therefore  $(B \setminus A) \cap D$  must be non-empty.

But if  $x \in B \setminus A$ , then  $x$  must be a limit point of  $A$ . But, as a consequence of theorem 5.5D if  $x$  is a limit point of  $A$  and if  $x$  lies in an open set  $D$ , then  $D \cap A \neq \emptyset$ . This contradicts the fact that  $A \subset C$  and  $C \cap D = \emptyset$ .

### Section 6.3

1. Suppose that  $B$  is a bounded subset of  $\mathbf{R}^2$ . Then there exists  $L$  such that  $B$  is contained inside the square  $S$  with corners (diametrically opposite)  $\langle -L, -L \rangle$  and  $\langle L, L \rangle$ . Let  $\varepsilon > 0$ .

Let  $h = \frac{\varepsilon}{\sqrt{2}}$ . Cover  $S$  with a rectangular grid of (closed) cells whose parallel sides are



distance  $h$  apart. Let  $N = \lceil \frac{2L}{h} \rceil + 1$ . Then, there are at most  $N^2$  cells in the grid, each cell has diameter  $\varepsilon$ , and the collection of cells in the grid covers  $S$  which contains  $B$ .

3. Example 1.  $S = \{s_n\}$  where each  $s_n = \langle 1/n, 0, 0, 0, 0, \dots \rangle$ , for  $n = 1, 2, 3, \dots$ . Then, each  $s_n \in \ell^2$  and  $\|s_n - s_m\|_2 = |1/n - 1/m|$ . It is easily seen that each subsequence of  $S$  is Cauchy.

Example 2.  $S = \{s_n\}$  where each  $s_n = \langle 0, 0, 0, \dots, 1/n, 0, 0, 0, \dots \rangle$ , where the single non-zero entry is in the  $n^{\text{th}}$  slot, for  $n = 1, 2, 3, \dots$ . Then, each  $s_n \in \ell^2$  and  $\|s_n - s_m\|_2 = (1/n^2 - 1/m^2)^{1/2}$ . It is easily seen that each subsequence of  $S$  is Cauchy.

4. Let  $S = \{x_1, x_2, \dots, x_n\}$  be a finite subset of  $M$ . Let  $\varepsilon > 0$ . Choose sets  $A_k = B(x_k, \varepsilon/2)$  for  $k = 1 \dots n$ . Then, the sets  $\{A_1, A_2, \dots, A_n\}$  cover  $S$  and each  $A_k$  has diameter  $\varepsilon$ .

5. Let  $\varepsilon > 0$ . Since  $M$  is total bounded there exist sets  $\{A_1, A_2, \dots, A_n\}$  in  $M$  such that  $\text{diam}(A_k) < \varepsilon$  and  $M \subset \bigcup_{k=1}^n A_k$ . But since  $A \subset M$ , we also have for  $B_k = A_k \cap A$  that

$$\text{diam}(B_k) < \varepsilon \text{ and } A \subset \bigcup_{k=1}^n B_k.$$

7. Denote  $A = A_0$ . We will show that there exists a sequence in  $A_0$  which is Cauchy. Since  $A_0$  is bounded there exists a closed bounded interval  $J_0$  such that  $A_0 \subset J_0$ . Let  $\text{diam}(J_0) = d$ .

Find the midpoint of  $J_0$  and divide  $J_0$  into two subintervals of equal width. Since  $A_0$  is infinite, at least one of these subintervals must contain an infinite number of points. Choose one of them and call it  $J_1$ . Note that  $\text{diam}(J_1) = d/2$ . Let  $A_1 = J_1 \cap A_0$ .

Find the midpoint of  $J_1$  and divide  $J_1$  into two subintervals of equal width. Since  $A_1$  is infinite, at least one of these subintervals must contain an infinite number of points. Choose one of them and call it  $J_2$ . Note that  $\text{diam}(J_2) = d/4$ . Let  $A_2 = J_2 \cap A_1$ .

Proceed inductively to find for each  $k = 1, 2, 3, \dots$  sets  $J_k$  such that each  $J_k \subset J_{k-1}$  and  $\text{diam}(J_k) = d/2^k$  and  $A_k = J_k \cap A_{k-1}$  where each  $A_k$  is infinite. The sets  $\{J_k\}$  satisfy the nested interval theorem and hence the property  $\bigcap_{k=1}^{\infty} J_k$  is a singleton.

Since each  $J_k \cap A_0 \neq \emptyset$  and  $\text{diam}(J_k) = d/2^k$  and  $\bigcap_{k=1}^{\infty} J_k \in J_k$ , then we see that  $\bigcap_{k=1}^{\infty} J_k$  is a cluster point of  $A_0$ .