

# Preliminary Examination 1998

## Complex Analysis

Do all problems.

Notation:

$$\mathbb{R} = \{ x : x \text{ is a real number} \}$$

$$B(a,r) = \{ z \in \mathbb{C} : |z - a| < r \}$$

$$\mathbb{C} = \{ z : z \text{ is a complex number} \}$$

$$\text{ann}(a,r_1,r_2) = \{ z \in \mathbb{C} : r_1 < |z - a| < r_2 \}$$

For  $G \subset \mathbb{C}$ , let  $\mathcal{A}(G)$  denote the set of analytic functions on  $G$  (mapping  $G$  to  $\mathbb{C}$ ).

1. Let  $f$  be an entire function.
  - (a) Suppose there exist  $a, b \in \mathbb{R}$  such that  $|f(z)| \leq (a\sqrt{|z|} + b)$  for all  $z \in \mathbb{C}$ . Show that  $f$  is constant.
  - (b) Suppose there exist  $a, b \in \mathbb{R}$  such that  $|f(z)| \leq (a|z|^{5/2} + b)$  for all  $z \in \mathbb{C}$ . What can you say about  $f$ ?
2. Let  $f, g$  be entire functions. Suppose  $|f(z)| \leq |g(z)|$  for all  $z \in \mathbb{C}$ . Prove there exists a constant  $c$  such that  $f \equiv cg$ .
3. Let  $A(r) = \text{ann}(0,r,1)$ ,  $0 < r < 1$ , and  $B = B(0,1) \setminus B(1/4,1/4)$ . Show that there exists an  $r$  such that  $A(r)$  is conformally equivalent to  $B$ .
4. Let  $G_1$  and  $G_2$  be simply connected regions, neither region is all of  $\mathbb{C}$ . Let  $a \in G_1$ . Suppose that  $f, g \in \mathcal{A}(G_1)$  such that  $f$  is one-to-one on  $G_1$  with  $f(G_1) = G_2$ ,  $g(G_1) \subset G_2$  and  $f(a) = g(a)$ . Prove  $|g'(a)| \leq |f'(a)|$ .
5. Define the Gamma function,  $\Gamma$ . Prove that  $\Gamma(z)\Gamma(1-z) = \pi/\sin(\pi z)$ , for  $z$  not an integer.
6. Let  $G$  be a region in  $\mathbb{C}$  and let  $\mathcal{F}$  be a subset of  $\mathcal{A}(G)$ . Prove that if  $\mathcal{F}$  is locally bounded, then  $\mathcal{F}$  is equicontinuous at each point of  $G$ .
7. Let  $D_1 = \{ z \in B(0,1) : \text{Im } z > 1/2 \}$ . Find a conformal map  $f$  which maps  $D_1$  one-to-one and onto  $B(0,1)$  such that  $f(3i/4) = 0$ .
8. Compute  $\int_{-\infty}^{\infty} \frac{e^{iax}}{(1+x^2)^2} dx, a > 0$ .
9. Find a non-constant function  $f \in \mathcal{A}(B(0,1))$  such that  $f$  has infinitely many zeros in  $B(0,1)$ .
10. Suppose  $\alpha \neq 0$  is a root of a polynomial  $p$  of degree  $n$  with rational coefficients. Prove that  $1/\alpha$  is a root of a polynomial of degree at most  $n$  with rational coefficients.